Original Research



Color Variability of Cosmopolitan Beetles in Mindanao, Philippines

Ernel D. Bagbag^{1*}), Mark Anthony J. Torres^{1,2}, Cesar G. Demayo^{1,3}

¹ Mindanao State University-Iligan Institute of Technology, Tibanga, Iligan City, Philippines

² School of Interdisciplinary Studies, MSU-Iligan Institute of Technology, Tibanga, Iligan City, Philippines

³ Center of Integrative Health, Premier Research Institute of Science and Mathematics, MSU-Iligan Institute of Technology, Tibanga, Iligan City, Philippines

Article Information

Received : February 20, 2024 Revised : March 10, 2024 Accepted : April 4, 2024 Published : April 10, 2024

Correspondence

Ernel D. Bagbag E-mail: ernel.bagbag@g.msuiit.edu.ph

Citation

D. Bagbag, E., Anthony J. M., & G. Demayo, C. (2024). Color Variability of Cosmopolitan Beetles in Mindanao, Philippines. *Andalasian International Journal of Entomology*, 2(1), 1–7. https://doi.org/10.25077/aijent.2.1.1-7.2024



Copyright: © 2024 by the authors. Submitted for possible **open access**

publication under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA) license

E-ISSN: <u>3026-2461</u> https://doi.org/10.25077/aijent.2.1.1-7.2024

INTRODUCTION

Color patterns in beetles, which result from the interplay of pigment and structural-based coloration, are pivotal in ensuring their survival. The enormous variety of this outstanding trait across Coleopterans has been shown to have evolved repeatedly, making it a remarkable trait to understand their diversification. Coloration patterns are a notable and accessible trait for study and experimental manipulation, consisting of multiple interacting components with measurable functions, making it an excellent model for

Abstract

Beetles (Insecta: Coleoptera) stands out as one of the most diverse insect groups, showcasing various color variations. These evolved color patterns present a fascinating trait crucial for understanding their evolution. However, studying these patterns poses challenges, given the intricate nature of colors in the natural world. While past researchers have explored beetle color patterns, quantifying colors requires costly equipment and sophisticated software. This paper introduces an alternative approach, utilizing digital images to examine color variability among Coleopterans. Forty-eight specimens from Mindanao, Philippines, were collected and photographed under consistent conditions. Subsequently, the images were calibrated and processed in R software to calculate image distances through cluster analysis. The results reveal that beetle color patterns fall into two categories: a dark color with deeper shades of gray and a moderately bright tone featuring a slight reddish hue with noticeable yellow highlights. Chrysochroa fulminans is the most distinct beetle across Coleopteran species due to its vibrant green coloration. Family-specific examination of color patterns revealed species with distinct color, Anomala flavipennis and A. smaragdina (Scarabaeidae), Otiorhynchus pauxillus (Curculionidae), Uloma culinaris (Tenebrionidae), and Nupserha fricator (Cerambycidae). The study's findings offer valuable insights into the evolution of Coleopterans, mainly their color patterns, serving as a valuable tool for classification.

Keywords

Digital images, cluster analysis, coleoptera, color variability, evolution

investigating evolutionary processes (Endler & Mappes, 2017). Moreover, colors provide information that may be used to identify and classify species, recognize individual characteristics, and indicate ecological or evolutionary elements of an organism's life (Badejo et al., 2020).

Beetles, constituting over 40% of all insect species, exhibit a vast diversity exceeding 450,000 species (Lawrence et al., 2011; McKenna et al., 2019; Sha'ari & Arumugam, 2019), with coloration patterns evolving to provide concealment from or deterrence of potential predators. These patterns serve various functions, including crypsis, masquerade, motion dazzle, motion camouflage, defense, signaling, and physiological adaptations (Stevens & Merilaita, 2008; Badejo et al., 2020).

The study of color patterns in organisms encompasses a broad range of ecological and evolutionary topics, such as plasticity, habitat choice, and color polymorphism (Boyle & Start, 2019; Stanbrook et al., 2021; Martín-Vega & Baz, 2013). However, categorizing Coleopterans based on color patterns is challenging due to the extensive diversity and intricacy of these patterns. Attempts to quantify colors often require expensive equipment. sophisticated software, and expertise (Van den Berg et al., 2020; Badiane et al., 2017; Weller & Westneat, 2019). Nonetheless, using digital cameras presents a more practical and efficient method for analyzing coloration in organisms (Kendal et al., 2013).

This paper adopts an alternative and practical approach to analyze beetle color patterns in the anthropogenic areas of Mindanao, Philippines. Moreover, this investigation provided a comprehensive overview of Coleopteran color patterns and compared them according to their family, genus, and species.

METHODS

Sampling Sites

The beetles were collected from eight anthropogenic areas of Mindanao, Philippines (Figure 1). Using opportunistic sampling techniques, pitfall entrapment, insect netting, and manual handpicking were imposed to collect beetle samples. Then, the specimen was placed in a container with 70% ethanol before being transported to the laboratory for digital profiling.



Figure 1. Map of the Collection Sites

Sample Preparation

Forty-eight (48) beetles, representing 12 different families, were gathered from various sampling sites included in the study. Table 1 provides the corresponding image codes assigned to each species, serving as labels for the subsequent graphs in this paper. The beetle samples were carefully positioned dorsally on a stereo microscope with a constantly aligned camera at a specific distance from the microscope eyepiece. Moreover, photographing the specimens took place in an enclosed dark room with exclusive illumination provided by the stereomicroscope, prohibiting other external light sources.

Table 1. Beetle species used in the study

Code	Species
SP01.jpg	Acalolepta rusticatrix
SP02.jpg	Aclees hirayamai
SP03.jpg	Anomala flavipennis
SP04.jpg	Anomala marginata
SP05.jpg	Anomala smaragdina
SP06.jpg	Apriona aphetor
SP07.jpg	Batocera magica
SP08.jpg	Carlschoenherria sulcipennis
SP09.jpg	Chalcosoma atlas
SP10.jpg	Cheilomenes sexmaculata
SP11.jpg	Chrysochroa fulminans
SP12.jpg	Cordylocera atricornis
SP13.jpg	Colophotia concolor
SP14.jpg	Cryptalaus lacteus
SP15.jpg	Cylindera discreta elaphroides
SP16.jpg	Cylindera minuta
SP17.jpg	Derosphaerus vicinus
SP18.jpg	Dorcus parvulus
SP19.jpg	Epepeotes plorator
SP20.jpg	Eretes griseus
SP21.jpg	Eucorynus crassicornis
SP22.jpg	Figulus sulcicollis
SP23.jpg	Holotrichia bipunctata
SP24.jpg	Hoplocerambyx spinicornis
SP25.jpg	Lema pectoralis
SP26.jpg	Leucopholis furforosa
SP27.jpg	Leucopholis pulverulenta
SP28.jpg	Macrolinus sulciperfectus
SP29.jpg	Metapocyrtus adspersus
SP30.jpg	Metriorrhynchus sp. (species 1)
SP31.jpg	Metriorrhynchus sp. (species 2)
SP32.jpg	Nupserha fricator
SP33.jpg	Onitis phartopus
SP34.jpg	Onthophagus hielkemai
SP35.jpg	Oryctes rhinoceros
SP36.jpg	Otiorhynchus pauxillus
SP37.jpg	Pentodon algerinus

SP38.jpg	Platymetopus flavilabris
SP39.jpg	Podontia quatuordecimpunctata
SP40.jpg	Prionocerus coeruleipennis
SP41.jpg	Prosoplus bankii
SP42.jpg	Protaetia fusca
SP43.jpg	Pseudozaena orientalis
SP44.jpg	Pterolophia crassipes
SP45.jpg	Serica sp.
SP46.jpg	Sybra ochreovittipennis
SP47.jpg	Uloma culinaris
SP48.jpg	Zophobas morio

Image Calibration

The digital images were processed using GIMP software version 2.13 (The GIMP Development Team, 2019). These images were calibrated through a simple white-balance correction and normalization. Moreover, the background of the photos was removed and replaced with a uniform white color. The resolution was adjusted to 700 pixels before saving the images as 24-bit Joint Photographic Experts Group (JPEG) files (Lenhert et al., 2011). These processed images were then organized into a separate folder for further analysis.

Colorimetric Analysis

Color quantification involves a three-step process: pixel binning, color conversion, and clustering (Figure 2). In pixel binning, the images were imported to R software (R Core Team, 2022), in which the plotPixels function of the colordistance package performed data reduction and normalization (Weller, 2021). In this process, the white background and photographic glares were excluded by setting the upper limit of RGB (Red, Green, and Blue) triplet colors to 0,0,0, the RGB triplet code for white color. This procedure eliminates unwanted elements from the images.

Then, the images were converted from RGB color space to Commission on Illumination 1976 L* a* b* (CIELab) using the LoadImage function. In the CIELab color space, Lightness (L*) is quantified on a scale from 0 (representing black) to 100 (representing white). Therefore, a higher L* value corresponds to a lighter color, while a lower value indicates a darker shade. The* channel spans from green to red on a scale of -120 to 120, where positive values represent red hues and negative values indicate green hues. On the other hand, the b* channel covers the range from blue to yellow on the same scale of -120 to 120, with positive values denoting yellow tones and negative values indicating blue tones (Ly et al., 2019).

A distance matrix was computed using the getColorDistanceMatrix function. This approach generated a heatmap, facilitating the comparison of each image based on the quantified color values. Then, the dominant color was determined by selecting the highest percentage of the quantified color that contributed to the image and then compared by beetle family, genus, and species.



Figure 2. Colorimetric Analysis Workflow

RESULTS AND DISCUSSION

Overview of Coleopteran Coloration

The clustered heatmap shows the overall color variability of Coleopterans sampled in this study, with blue cells indicating a high degree of similarity (low color distance scores) and pink cells showing low similarity (high color distance scores). Based on the cluster heatmap, the color patterns of the Coleopterans collected from the anthropogenic environments are divided into two distinct clusters (Figure 3-A).





Figure 3. Color variability of Coleopterans (A). Overview of Coleopteran Coloration, (B) Clustering of Coleopterans based on their Body Coloration (Images not drawn to scale)

The first cluster is composed of darker beetles, with colors ranging from gray to black, while the second group is characterized by light-colored beetles ranging from red to yellow and orange. In the first cluster, the species A. smaragdina and C. fulminans are the most distinct beetles in this group, showing a distinctive greenish coloration. Furthermore, C. fulminans is the species that is distinct across Coleopteran species, as indicated by low similarity scores (pink cells). On the other hand, L. pectoralis, Ρ. quatuordecimpunctata, Metriorrhynchus SD. (species 2), and N. fricator are beetles with distinct colors compared to the first cluster. The beetles in their respective clusters are depicted in Figure 3-B.

Color Variability Between Species Within Family

A comparative analysis of beetle species within their families was conducted to explore color variations. The Scarabaeidae, Curculionidae, Tenebrionidae, and Cerambycidae were specifically analyzed, as these families had more than three species in the study. In the Scarabaeidae family, the results identified A. flavipennis and A. smaragdina as distinct species within the Scarab samples. Another group of scarab beetles exhibited darker colors, featuring prominent dark gray to red elements in their bodies in contrast to A. flavipennis and A. smaragdina (Figure 4-A). Within the Curculionidae family (weevils), O. pauxillus stood out as distinct from other weevils in this study, displaying a reddish coloration with a hint of yellow. Conversely, M. adspersus and A. hirayamai formed a group with darker colors than O. pauxillus (Figure 4-B). U. culinaris emerged as a distinct species among Tenebrionidae (darkling beetles) when colors were compared with D. vicinus and Z. morio (Figure 4-C). Furthermore, N. fricator emerged as a distinct Cerambycidae beetle species (Longhorn beetles) among the ten beetles sampled in the present study (Figure 4-D).





Figure 4. Color Variation of Beetle Species Based on Their Family Groups. (A) Scarabaeidae, (B) Curculionidae, (C) Tenebrionidae, (D) Cerambycidae (images are not drawn to scale)

Discussion

This paper presents a comprehensive classification of bettle coloration, highlighting two distinct categories: (1) deeper shades of gray and dark colors and (2) moderately bright tones with a slight reddish hue and noticeable yellow highlights. Moreover, the cluster heatmap revealed the occurrence of analogous individuals among Coleoptera, which denotes shared coloration patterns. The diverse coloration patterns observed and parallelism among Coleopterans are ascribed more to their functional morphology and evolutionary significance than their taxonomic position.

The emergence of these color patterns can be attributed to the presence of melanins, encompassing a spectrum ranging from deep black to rich reddishbrown, as well as the contributions of pterins, ommochromes, and carotenoids that infuse vibrant red, orange, and yellow hues (Badejo et al., 2020). These compounds can be transferred to the beetles through ingestion, causing them to express body coloration.

In leaf beetles (Chrysomelidae), those with a diverse range of host plants generally exhibit reduced red coloration compared to their counterparts with a more specialized diet. This implies that beetles with versatile dietary preferences may adopt a subdued red hue as a strategic measure to lower their visibility while navigating different ecological environments (Tan et al., 2017). This observation aligns with the coloration of the studied Chrysomelid beetles, *L. pectoralis* and *P. quatuordecimpunctata*, both recognized as polyphagous species, displaying a subtle red-orange tint in their bodies.

Interestingly, the polyphagous Melolonthinae beetles (C. sulcipennis and H. bipunctata) and A. flavipennis exhibit a prominent red element in their bodies. This coloration pattern is likely attributed to their shared dietary source, as insects obtain carotenoids (the pigments responsible for red, orange, and yellow hues) from their food (Badejo et al., 2020). These beetles gain a significant advantage from this coloration strategy, as the combination of red, orange, yellow, and white hues contributes to aposematism (Badejo et al., 2020). In the case of the ladybird Asian beetle Harmonia axvridis (Coccinellidae), the presence of red-orange elytra coloration indicates the conveyance of details regarding chemical defense to potential mates or predators (Bezzerides et al., 2007).

The diversity in color patterns among beetles is also attributed to their aposematic functions. Badejo and colleagues (2020) underscored that aposematic coloration reduces predation risk. For instance, Geotrupes beetles, G. auratus and G. laevistriatus, have evolved similar bright colorations, acting as a warning signal, especially for visually-oriented predators (Watanabe et al., 2022). Additionally, color patterns hold evolutionary significance for thermoregulation (Badejo et al., 2020). Insects with darker colors gain enhanced protection against ultraviolet radiation and excessive heat (True, 2003). For example, in *Harpalus affinis* (Carabidae), a dark bronze shade conferred selective advantages during periods of heightened pollution in urban areas, suggesting that varying degrees of urbanization could influence the prevalence of distinct color patterns in beetles (Keinath et al., 2020).

This study suggests that beetle coloration has undergone evolutionary changes over time, with these body colorations holding distinct evolutionary significance for the survival of beetles. Moreover, the observed variability in body coloration among Coleopterans indicates that species classification cannot rely solely on a single trait but requires the consideration of multiple characters.

CONCLUSIONS

This study employed digital images to explore the coloration patterns of beetles in Mindanao, Philippines. The analysis yielded a comprehensive examination of Coleopteran color variability, contributing to our understanding of the evolutionary processes that led to their diversification. The research results suggest that specific Coleopteran groups can be effectively classified through

quantitative color analysis, presenting a potential tool for the systematic classification of Coleopterans.

Utilizing digital image-based color profiling presents a non-invasive methodology; however, it is imperative to acknowledge that some beetle species exhibit iridescent color patterns. These patterns can considerably influence the results, particularly when observing the beetle from various angles. Furthermore, it is also essential to delve into the underlying color mechanisms to investigate how distinct non-homologous structures can contribute to chromatic effects. Recognizing this limitation and addressing it in subsequent research endeavors is imperative.

ACKNOWLEDGMENT

The authors would like to acknowledge the Philippine Department of Science and Technology under the Accelerated Science and Technology Human Resource Development Program for the research grant.

REFERENCES

Badejo, O., Skaldina, O., Gilev, A., & Sorvari, J. (2020). Benefits of insect colours: a review from social insect studies. *Oecologia*, *194*, 27-40. https://doi.org/10.1007/s00442-020-04738-1.

Badiane, A., de Lanuza, G. P., Custodio, M., & Carazo, P. (2017). Colour patch size and measurement error using reflectance spectrophotometry. *Methods in Ecology and Evolutipm, 8*, 1585-1593. https://doi.org/10.1111/204 1-210X.12801.

Bezzerides, A. L., McGraw, K., Parker, R. S., & Husseini, J. (2007). Elytra color as a signal of chemical defense in the Asian ladybird beetle Harmonia axyridis. *Behavioral Ecology and Sociobiology*, *61*, 1401–1408. https://doi.org/10.100 7/s00265-007-0371-9.

Boyle, J., & Start, D. (2019). Plasticity and habitat choice match colour to function in an ambush bug. *Functional Ecology*, *34*, 822-829. https://doi.org/ 10.1111/1365-2435.13528.

Endler, J. A., & Mappes, J. (2017). The current and future state of animal coloration research. *Philosophical Transactions of the Royal Society B, 372* (1724). https://doi.org/10.1098/rstb.2016.0352.

Keinath, S., Frisch, J., Müller, J., Mayer, F., & Rödel, M.-O. (2020). Spatio-Temporal Color Differences Between Urban and Rural Populations of a Ground Beetle During the Last 100 Years. *Frontiers in* *Ecology and Evolution, 7.* https://doi.org/10.3389/fe vo.2019.00525.

Kendal, D., Hausser, C. E., Garrad, G. E., Jellinek, S., Gilijohann, K. M., & Moore, J. L. (2013). Quantifying Plant Colour and Colour Difference as Perceived by Humans Using Digital Images. *PLoS ONE, 8*(8). https://doi.org/10.1371/journal.pone.0072 296.

Lawrence, J. F., Slipinski, A., Seago, A., Thayer, M., Newton, A., & Marvaldi, A. (2011). Phylogeny of the Coleoptera based on Morphological Characters of Adults and Larvae. *Annales Zoologici, 61*(1), 1-217. https://doi.org/10.3161/000345411X576725.

Lenhert, M., Balaban, M., & Emmel, T. C. (2011). A New Method for Quantifying Color of Insects. *Florida Entomologist, 94*(2), 201-207. https://doi.org/10.165 3/024.094.0212.

Ly, B. K., Dyer, E. B., Feig, J. L., Chien, A. L., & Del Bino, S. (2019). Research Techniques Made Simple: CutaneousColorimetry: A Reliable Technique for ObjectiveSkin Color Measurement. *Journal of Investigative Dermatology*, *140*(1), 3-12. https://doi.org/10.1016/j.jid.2019.11.003.

Martín-Vega, D., & Baz, A. (2013). Variation in the colour of the necrophagous fly, Prochyliza nigrimana (Diptera: Piophilidae): A case of seasonal polymorphism. *European Journal of Entomology, 108*(2), 231-234. https://doi.org/10.14411/eje. 2011.031.

McKenna, D., Shin, S., Ahrens, D., Balke, M., Beza-Beza, C., Clarke, D. J., . . . Beutel, R. (2019). The evolution and genomic basis of beetle diversity. *Proceedings of the National Academy of Science*, *116*(49), 24729-24737. https://doi.org/10.1073/pnas. 1909655116.

R Core Team. (2022). R: A language and environment for statistical computing. Vienna, Austrria: R Foundation for Statistical Computing. https://www.R-project.org/.

Sha'ari, N., & Arumugam, N. (2019). Beetles Diversity (Order: Coleoptera) of R.E.A.C.H. Biodiversity Center, Cameron Highlands, Pahang, Malaysia. *Serangga, 24*(2), 58-67.

Stanbrook, R. A., Harris, W. E., Wheater, C. P., & Jones, M. (2021). Evidence of phenotypic plasticity along an altitudinal gradient in the dung beetle Onthophagus proteus. *PeerJ*, *9*, e10798. https://doi.org/10.7717/peerj.10798.

Stevens, M., & Merilaita, S. (2008). Animal camouflage: current issues and new perspectives. *Philosophical Transactions of the Royal Society B,*

364, 423–427. https://doi.org/10.1098/rstb.2008 .0217.

Tan, E. J., Reid, C. A., Symonds, M. R., Jurado-Rivera, J., & Elgar, M. A. (2017). The Role of Life-History and Ecology in the Evolution of Color Patterns in Australian Chrysomeline Beetles. *Frontiers in Ecology and Evolution*, *5*, 140. https://doi/org/10.3389/fevo.2017 .00140.

The GIMP Development Team. (2019). GIMP. https://www.gimp.org.

True, J. R. (2003). Insect melanism: the molecules matter. *Trends in Ecology and Evolution, 18*(12), 640-647. https://doi.org/10.1016/j.tree.2003.09.006

Van den Berg, C. P., Troscianko, J., Endler, J. A., Marshall, N. J., & Cheney, K. L. (2020). Quantitative Colour Pattern Analysis (QCPA): A Comprehensive framework for the analysis of colour patterns in nature. *Methods in Ecology and Evolution, 11*, 316-332. https://doi.org/10.1111/2041-210X.13328.

Watanabe, T., Tanigaki, T., Nishi, H., Ushimaru, A., & Takeuchi, T. (2002). A Quantitative Analysis of Geographic Color Variation in Two *Geotrupes* Dung Beetles. *Zoological Science*, 351-358. https://doi.org/10.2108/zsj.19.351.

Weller, H. I., & Westneat, M. W. (2019). Quantitative color profiling of digital images with earth mover's distance using the R package colordistance. *PeerJ*, 7. https://doi.org/10.7717/peerj.6398.